

# TRANSLATING THE CANTOR SET BY A RANDOM

RANDALL DOUGHERTY, JACK LUTZ, R. DANIEL MAULDIN, AND JASON TEUTSCH

ABSTRACT. We determine the constructive dimension of points in random translates of the Cantor set. The Cantor set “cancels randomness” in the sense that some of its members, when added to Martin-Löf random reals, identify a point with lower constructive dimension than the random itself. In particular, we find the Hausdorff dimension of the set of points in a Cantor set translate with a given constructive dimension.

## 1. FRACTALS AND RANDOMS

We explore an essential interaction between algorithmic randomness, classical fractal geometry, and additive number theory. In this paper, we consider the dimension of the intersection of a given set with a translate of another given set. We shall concern ourselves not only with classical Hausdorff measures and dimension but also the effective analogs of these concepts.

More specifically, let  $\mathcal{C}$  denote the standard middle third Cantor set [6, 18], and for each number  $\alpha$  let

$$(1.1) \quad \mathcal{E}_{=\alpha} = \{x : \text{cdim}_{\mathbb{H}}\{x\} = \alpha\}$$

consist of all real numbers with constructive dimension  $\alpha$ . We answer a question posed to us by Doug Hardin by proving the following theorem: if  $1 - \log 2 / \log 3 \leq \alpha \leq 1$  and  $r$  is a Martin-Löf random real, then the Hausdorff dimension of

$$(1.2) \quad (\mathcal{C} + r) \cap \mathcal{E}_{=\alpha}$$

is  $\alpha - (1 - \log 2 / \log 3)$ . From this result we obtain a simple relation between the effective and classical Hausdorff dimensions of (1.2); the difference is exactly 1 minus the dimension of the Cantor set. We conclude that many points in the Cantor set additively cancel randomness.

We discuss some of the notions involved in this paper. Intuitively, a real is “random” if it does not inherit any special properties by belonging to an effective null class. We say a number is *Martin-Löf random* [3, 12] if it “passes” all Martin-Löf tests. A *Martin-Löf test* is a uniformly computably enumerable (c.e.) sequence [3, 16] of open sets  $\{U_m\}_{m \in \mathbb{N}}$  with  $\lambda(U_m) \leq 2^{-m}$ , where  $\lambda$  denotes Lebesgue measure [18]. A number  $x$  *passes* such a test if  $x \notin \bigcap_m U_m$ .

The *Kolmogorov complexity* of a string  $\sigma$ , denoted  $K(\sigma)$ , is the length (in this paper we will measure length in ternary units) of the shortest program (under a fixed universal machine) which outputs  $\sigma$  [8]. For a real number  $x$ ,  $x \upharpoonright n$  denotes the

---

2000 *Mathematics Subject Classification.* Primary 68Q30; Secondary 11K55, 28A78.

*Key words and phrases.* algorithmic randomness, fractal geometry, additive number theory.

Research supported by NSF Grants 0652569 and 0728806.

Research supported by NSF Grant DMS-0700831.

Research supported by Deutsche Forschungsgemeinschaft grant ME 1806/3-1.

first  $n$  digits in a ternary expansion of  $x$ . Martin-Löf random reals have high initial segment complexity [3]; indeed every Martin-Löf random real  $r$  satisfies  $\lim_n K(r \upharpoonright n)/n = 1$ . This fact conforms with our intuition that random objects do not compress much.

We introduce a couple of classical dimension notions. Let  $E \subseteq \mathbf{R}^n$ . The *diameter* of  $E$ , denoted  $|E|$ , is the maximum distance between any two points in  $E$ . We will use  $\text{card}$  for cardinality. A *cover*  $\mathcal{G}$  for a set  $E$  is a collection of sets whose union contains  $E$ , and  $\mathcal{G}$  is a  $\delta$ -*mesh cover* if the diameter of each member  $G$  is at most  $\delta$ . For a number  $\beta \geq 0$ , the  $\beta$ -*dimensional Hausdorff measure* of  $E$ , written  $\mathcal{H}^\beta(E)$ , is given by  $\lim_{\delta \rightarrow 0} \mathcal{H}_\delta^\beta(E)$  where

$$(1.3) \quad \mathcal{H}_\delta^\beta(E) = \inf \left\{ \sum_{G \in \mathcal{G}} |G|^\beta : \mathcal{G} \text{ is a countable } \delta\text{-mesh cover of } E \right\}.$$

The *Hausdorff dimension* of a set  $E$ , denoted  $\dim_{\text{H}}(E)$ , is the unique number  $\alpha$  where the  $\alpha$ -dimensional Hausdorff measure of  $E$  transitions from being negligible to being infinitely large; if  $\beta < \alpha$ , then  $\mathcal{H}^\beta(E) = \infty$  and if  $\beta > \alpha$ , then  $\mathcal{H}^\beta(E) = 0$  [6, 18]. Let  $S_\delta(E)$  denote the smallest number of sets of diameter at most  $\delta$  which can cover  $E$ . The *upper box-counting dimension* [6] of  $E$  is defined as

$$\overline{\dim}_{\text{B}}(E) = \limsup_{\delta \rightarrow 0} \frac{\log S_\delta(E)}{-\log \delta}.$$

The *effective* (or *constructive*)  $\beta$ -*dimensional Hausdorff measure* of a set  $E$ ,  $\mathfrak{c}\mathcal{H}^\beta(E)$ , is defined exactly in the same way as Hausdorff measure with the restriction that the covers be uniformly c.e. open sets [3, Definition 13.3.3]. This yields the corresponding notion of the *effective* (or *constructive*) *Hausdorff dimension* of a set  $E$ ,  $\text{cdim}_{\text{H}} E$ . Lutz [10] showed

$$(1.4) \quad \text{cdim}_{\text{H}} E = \sup \{ \text{cdim}_{\text{H}} \{x\} : x \in E \},$$

and from work of Mayordomo [14]( $\geq$ ) and Levin [7]( $\leq$ ) (also see [3]) we have for any real number  $x$ ,

$$(1.5) \quad \text{cdim}_{\text{H}} \{x\} = \liminf_{n \rightarrow \infty} \frac{K(x \upharpoonright n)}{n}.$$

Mayordomo and Levin prove their results in  $\{0, 1\}^{\mathbf{N}}$ , but the results carry over to the reals. We define the *constructive dimension* of a point  $x$  to be the effective Hausdorff dimension of the singleton  $\{x\}$ . A further effective dimension notion, the *effective packing dimension* [1, 3] satisfies

- $\text{cdim}_{\text{P}} \{x\} = \limsup_{n \rightarrow \infty} \frac{K(x \upharpoonright n)}{n}$ , and
- $\text{cdim}_{\text{P}} \mathcal{C} = \dim_{\text{H}} \mathcal{C}$ .

Let us first note some simple bounds on the complexity of a point in the translated Cantor set  $\mathcal{C} + r = \{y : (\exists x \in \mathcal{C}) [y = x + r]\}$ .

**Theorem 1.1.** *Let  $x \in \mathcal{C}$  and let  $r$  be a Martin-Löf random real. Then*

$$1 - \dim_{\text{H}} \mathcal{C} \leq \text{cdim}_{\text{H}} \{x + r\} \leq 1.$$

*Proof.* Given  $x$  and  $r$ , there is a constant  $c$  such that for all  $n$ ,

$$K[(x + r) \upharpoonright n] + K(x \upharpoonright n) \geq K(r \upharpoonright n) + c.$$

Thus,

$$\begin{aligned} 1 \geq \text{cdim}_{\mathbb{H}}\{x+r\} &= \liminf_{n \rightarrow \infty} \frac{K[(x+r) \upharpoonright n]}{n} \\ &\geq \liminf_{n \rightarrow \infty} \frac{K(r \upharpoonright n)}{n} - \limsup_{n \rightarrow \infty} \frac{K(x \upharpoonright n)}{n} \\ &\geq 1 - \text{cdim}_{\mathbb{P}}\{x\} \geq 1 - \text{cdim}_{\mathbb{P}}\mathcal{C} = 1 - \dim_{\mathbb{H}}\mathcal{C}. \quad \square \end{aligned}$$

In Section 2, we will indicate how some points cancel randomness. We show that for every  $r$  there exists an  $x$  such that the constructive dimension of  $x+r$  is as close to the lower bound as one likes. Later we will show that for each  $r$  and number  $\alpha$  within the correct bounds, not only does there exist some  $x \in \mathcal{C}$  so that  $x+r$  has constructive dimension  $\alpha$ , but we will determine the Hausdorff dimension of the set of all  $x$ 's with constructive dimension  $\alpha$ . At this point let us give a heuristic argument indicating what the Hausdorff dimension of this set might be.

Fix a number  $1 - \dim_{\mathbb{H}}\mathcal{C} < \alpha < 1$ , and following the notation in (1.1), let

$$\mathcal{E}_{\leq \alpha} = \{x : \text{cdim}_{\mathbb{H}}\{x\} \leq \alpha\}.$$

From [10] (see also [2]), we know that the effective Hausdorff dimension of  $\mathcal{E}_{\leq \alpha}$  satisfies  $\text{cdim}_{\mathbb{H}}\mathcal{E}_{\leq \alpha} = \dim_{\mathbb{H}}\mathcal{E}_{\leq \alpha} = \alpha$ . Since the upper box counting dimension of  $\mathcal{C}$  satisfies  $\overline{\dim}_{\mathbb{B}}\mathcal{C} = \dim_{\mathbb{H}}\mathcal{C}$  [6, Example 3.3], we have  $\dim_{\mathbb{H}}(\mathcal{C} \times \mathcal{E}_{\leq \alpha}) = \dim_{\mathbb{H}}\mathcal{C} + \alpha$  [6, Corollary 7.4]. Define  $f : \mathbf{R}^2 \mapsto \mathbf{R}^2$  by

$$(1.6) \quad f(x, y) = (y - x, y).$$

Then  $f$  is a bi-Lipschitz map and therefore preserves Hausdorff dimension [5, Lemma 1.8]. So, letting  $B = f^{-1}(\mathcal{C} \times \mathcal{E}_{\leq \alpha})$ , we have  $\dim_{\mathbb{H}}B = \dim_{\mathbb{H}}\mathcal{C} + \alpha$ . The *vertical fiber* of  $B$  at  $x$ , or set of points  $y$  such that  $(x, y) \in B$ , is

$$(1.7) \quad B_x = (\mathcal{C} + x) \cap \mathcal{E}_{\leq \alpha}.$$

Let  $\gamma > \dim_{\mathbb{H}}\mathcal{C} + \alpha$ . By the Fubini type inequality for Hausdorff measures [5, Theorem 5.12], there is a positive constant  $b$  such that

$$0 = \mathcal{H}^{\gamma}(B) \geq b \int \mathcal{H}^{\gamma-1}(B_x) d\mathcal{H}^1(x) = b \int \mathcal{H}^{\gamma-1}(B_x) dx.$$

So for Lebesgue measure a.e.  $x$ ,  $\mathcal{H}^{\gamma-1}[(\mathcal{C} + x) \cap \mathcal{E}_{\leq \alpha}] = 0$ . Therefore, for Lebesgue measure a.e.  $x$ ,

$$\dim_{\mathbb{H}}[(\mathcal{C} + x) \cap \mathcal{E}_{\leq \alpha}] \leq \alpha - (1 - \dim_{\mathbb{H}}\mathcal{C}).$$

We would like to turn this inequality into an equality for every Martin-Löf random real  $x$ , but even showing that inequality holds for all Martin-Löf randoms is a problem. This is because, in general, if one has a non-negative Borel measurable function  $f$  and  $\int f(x) dx = 0$ , then  $f(x) = 0$  for Lebesgue measure almost every  $x$ , but there may be Martin-Löf random  $x$ 's for which  $f(x) > 0$ . In Section 4 of this paper, we more carefully analyze our particular situation to obtain the conjectured upper bound.

## 2. SOME POINTS CANCEL RANDOMNESS

We begin with a simple example illustrating how points in the Cantor set can counteract randomness. Let us briefly review some facts about the standard middle-third Cantor set.

(1) We may express any  $x \in [0, 1]$  as a ternary expansion:

$$x = .x_1x_2x_3\dots = \sum_{n=1}^{\infty} \frac{x_n}{3^n}$$

where each  $x_n \in \{0, 1, 2\}$ . The Cantor set  $\mathcal{C}$  consists of those  $x$  for which the  $x_n$ 's are all 0 or 2, and the half-size Cantor set  $\frac{1}{2}\mathcal{C}$  consists of those  $x$  for which the  $x_n$ 's are all 0 or 1.

- (2) Any number in the interval  $[0, 2]$  can be written as a sum of two elements of the Cantor set. Indeed  $\frac{1}{2}\mathcal{C} + \frac{1}{2}\mathcal{C} = [0, 1]$  because the coordinates of any ternary decimal can be written as  $0 + 0$ ,  $0 + 1$ , or  $1 + 1$ .
- (3) The Hausdorff dimension and effective Hausdorff dimension of the Cantor set agree (see [5, Theorem 1.4] and [15, Section 1.7.1]):

$$\dim_{\mathbb{H}} \mathcal{C} = \text{cdim}_{\mathbb{H}} \mathcal{C} = \frac{\log 2}{\log 3} \approx 0.6309.$$

All the usual notions of dimension: Hausdorff, packing, upper and lower Minkowski or box counting, agree on  $\mathcal{C}$  [6].

Since the Cantor set contains the point 0, it is immediate that  $\mathcal{C} + r$  contains points of constructive dimension 1 whenever  $r$  is Martin-Löf random. We now present a simple construction which identifies points with lower constructive dimension.

**2.1. A point within 2/3 of optimal.** Let  $r \in [0, 1]$  be a real with ternary expansion  $.r_1r_2\dots$ . Choose  $t = .t_1t_2\dots \in \mathcal{C}$  as follows. Let

$$t_n = \begin{cases} 0 & \text{if } r_n \in \{1, 2\}, \\ 2 & \text{otherwise.} \end{cases}$$

Then

$$r_n + t_n = \begin{cases} 2 & \text{if } r_n = 0, \\ 1 & \text{if } r_n = 1, \\ 2 & \text{if } r_n = 2, \end{cases}$$

Since  $(0, \frac{1}{3}, \frac{2}{3})$  is the limiting frequency probability vector for this sequence, the constructive dimension of this sequence is dominated by the effective Hausdorff dimension of the set of all sequences with this limiting frequency vector. By [10, Lemma 7.3], we have

$$\text{cdim}_{\mathbb{H}}\{r + t\} \leq \text{entropy} \left(0, \frac{1}{3}, \frac{2}{3}\right) = -\frac{1}{3} \log_3 \frac{1}{3} - \frac{2}{3} \log_3 \frac{2}{3} = 1 - \frac{2}{3} \cdot \dim_{\mathbb{H}} \mathcal{C}.$$

This shows that for every  $r$ , there exists some point in  $\mathcal{C} + r$  whose constructive dimension is at most  $1 - (2/3) \dim_{\mathbb{H}} \mathcal{C}$ . Next we construct points whose constructive dimensions approach the  $1 - \dim_{\mathbb{H}} \mathcal{C}$  limit given in Theorem 1.1.

**2.2. Building blocks: achieving near the limit.** We consider a more refined example. Recall that  $\frac{1}{2}\mathcal{C}$  is the set of all ternary decimals in  $[0, 1]$  made from 0's and 1's (and no 2's), and take  $\frac{1}{2}E_3$  to be the set of ternary decimals in  $[0, 1]$  generated from concatenated blocks in

$$B_3 = \{000, 002, 021, 110, 112\}.$$

So in particular  $\frac{1}{2}\mathcal{C}$  is generated by concatenating the blocks

$$(2.1) \quad C_3 = \{000, 001, 010, 011, 100, 101, 110, 111\}.$$

By exhaustion, any ternary block of length 3 can be written as the sum of a member of  $C_3$  plus a member of  $B_3$  (e.g.  $020 = 002 + 011$ ). Therefore  $\frac{1}{2}\mathcal{C} + \frac{1}{2}E_3 = [0, 1]$ , and furthermore, as we shall see in (2.3),

$$\text{cdim}_{\mathbf{H}} E_3 \leq \frac{\log 5}{\log 27} \approx 0.488324507.$$

The following are examples of optimal complementary block sets for each length (in terms of size). These blocks are not unique: for each length  $k$ , there is more than one smallest block set which can be added to the length  $k$  analogue of (2.1) in order to achieve all ternary numbers up to length  $k$ .

$$\begin{aligned} B_1 &= \{0, 1\}, \\ B_2 &= \{00, 02, 11\}, \\ B_3 &= \{000, 002, 021, 110, 112\}, \\ B_4 &= \{0000, 0002, 0011, 0200, 0202, 0211, 1100, 1102, 1111\}, \\ B_5 &= \{00000, 00002, 00021, 00112, 00210, 01221, 02012, \\ &\quad 02110, 02201, 10212, 11010, 11101, 11120, 11122\}. \end{aligned}$$

Note that  $B_4$  is just the product  $B_2 \times B_2$  and is still optimal. We wonder whether products can be optimal for larger indices as well.

A set  $E \subseteq \mathbf{R}$  is called *computably closed* if there exists a computable predicate  $R$  such that  $x \in E \iff (\forall n) R(x \upharpoonright n)$ . We shall use the following combinatorial lemma of Lorentz to prove that there exist sufficiently small complementary blocks for each length whose members can be concatenated to achieve computably closed sets with low effective Hausdorff dimension (Theorem 2.1).

**Lorentz's Lemma** ([9]). *There exists a constant  $c$  such that for any integer  $k$ , if  $A \subseteq [0, k]$  is a set of integers with  $|A| \geq \ell \geq 2$ , then there exists a set of integers  $B \subseteq (-k, k)$  such that  $A + B \supseteq [0, k]$  with  $|B| \leq ck \frac{\log \ell}{\ell}$ .*

Although Lorentz's Lemma as such does not appear explicitly in Lorentz's original paper, as mentioned in [4], his argument in [9, Theorem 1] proves the statement above.

**Theorem 2.1.** *There exists a uniform sequence of computably closed sets  $E_1, E_2, \dots$  such that*

- (I)  $\frac{1}{2}\mathcal{C} + \frac{1}{2}E_n = [0, 1]$  for all  $n$ , and
- (II)  $\lim_{n \rightarrow \infty} \text{cdim}_{\mathbf{H}} E_n = 1 - \dim_{\mathbf{H}} \mathcal{C}$ .

*Proof.* For  $k > 0$ , let

$$I_k = \{i : 0 \leq i < 3^k\},$$

and let

$$\begin{aligned} C_k &= \{i : 0 \leq i < 3^k \text{ and } i \text{ has only 0's and 1's in its ternary expansion}\} \\ &= \left\{ \sum_{j=0}^{k-1} \delta_j 3^j : \delta_j \in \{0, 1\} \right\}. \end{aligned}$$

By Lorentz's Lemma (applied to  $C_k$ ), there exists a set  $B_k \subseteq (-3^k, 3^k)$  with

$$(2.2) \quad I_k \subseteq C_k + B_k$$

satisfying

$$\text{card } B_k \leq c' \cdot 3^k \cdot \frac{\log(2^k)}{2^k} = c' k \log 2 \left(\frac{3}{2}\right)^k = ck \left(\frac{3}{2}\right)^k,$$

where  $c'$  is the constant from Lorentz's Lemma and  $c = c' \log 2$ . Set

$$\frac{1}{2}\mathcal{C} = \left\{ \sum_{n=1}^{\infty} \frac{a_n}{3^{kn}} : a_n \in C_k \right\} \quad \text{and} \quad \frac{1}{2}E_k = \left\{ \sum_{n=1}^{\infty} \frac{b_n}{3^{kn}} : b_n \in B_k \right\}.$$

Let  $x \in [0, 1]$  have ternary expansion

$$0.x_1x_2x_3\dots = \sum_{n=1}^{\infty} \sum_{j=1}^k \frac{x_{(n-1)k+j}}{3^{(n-1)k+j}} = \sum_{n=1}^{\infty} \sum_{s=0}^{k-1} \frac{x_{nk-s}}{3^{nk-s}} = \sum_{n=1}^{\infty} \frac{1}{3^{nk}} \left( \sum_{s=0}^{k-1} x_{nk-s} 3^s \right).$$

By (2.2), there exist sequences  $\{a_n\}$  with members in  $C_k$  and  $\{b_n\}$  from  $B_k$  such that for all  $n \geq 1$ ,

$$\sum_{s=0}^{k-1} x_{nk-s} 3^s = a_n + b_n,$$

and therefore

$$x = \sum_{n=1}^{\infty} \frac{a_n}{3^{kn}} + \sum_{n=0}^{\infty} \frac{b_n}{3^{kn}} \in \frac{1}{2}\mathcal{C} + \frac{1}{2}E_k,$$

which proves part (I).

Define

$$\gamma_k = \frac{\log(\text{card } B_k)}{\log 3^k} \leq 1 - \frac{\log 2}{\log 3} + \frac{\log c + \log k}{k \log 3}.$$

To prove part (II), we first note that  $\text{cdim}_{\mathbb{H}}(E_k) \leq \gamma_k$ . For every  $n > 0$ , we can uniformly cover  $E_k$  with  $(\text{card } B_k)^n$  intervals of size  $3 \cdot 3^{-kn}$ . Indeed, there are  $\text{card } B_k$  choices for each of the first  $n$  blocks in any member of  $E_k$ , and a closed interval of length  $3 \cdot 3^{-kn}$  covers all possible extensions of each such prefix. Each  $E_k$  is a computably closed set and we have:

$$(2.3) \quad c\mathcal{H}^{\gamma_k}(E_k) \leq \lim_{n \rightarrow \infty} (\text{card } B_k)^n \cdot 3^{\gamma_k} \cdot (3^{-kn})^{\gamma_k} = 3^{\gamma_k}.$$

So,  $\limsup_{k \rightarrow \infty} \dim_{\mathbb{H}} E_k \leq 1 - \dim_{\mathbb{H}} \mathcal{C}$ . Also, we have  $\gamma_k \geq \overline{\dim}_{\mathbb{B}}(E_k)$ . Again, applying the fact the Lipschitz map  $(x, y) \mapsto x + y$  doesn't increase dimension [5, Lemma 1.8] together with a bound on the dimension of a product set in terms of the dimension of its factors ([6, Product formula 7.3]) we have

$$(2.4) \quad 1 = \dim_{\mathbb{H}}(\mathcal{C} + E_k) \leq \dim_{\mathbb{H}}(\mathcal{C} \times E_k) \leq \overline{\dim}_{\mathbb{B}} \mathcal{C} + \dim_{\mathbb{H}} E_k = \dim_{\mathbb{H}} \mathcal{C} + \dim_{\mathbb{H}} E_k.$$

The leftmost equality of (2.4) follows from part (I) and the rightmost equality follows from [6, Example 3.3]. Thus part (II) holds.  $\square$

From the construction of the set  $E_k$  one would think that  $\dim_{\mathbb{H}} E_k = \gamma_k$  and  $0 < \mathcal{H}^{\gamma_k}(E_k) < \infty$ . However, it is not clear that the similarity maps that one might naturally use to generate the self-similar set  $E_k$  satisfy the open set condition, see [5]. In fact, there are possible cases (e.g., when  $B_k$  contains two consecutive numbers and two numbers that differ by  $3^k$ ) where we would get  $\dim_{\mathbb{H}} E_k < \gamma_k$ .

We obtain immediately from Theorem 2.1 the following:

**Corollary 2.2.** *For every real  $r \in [0, 2]$  and every  $\epsilon > 0$ , there exists a point in  $\mathcal{C} + r$  whose constructive dimension is less than  $1 - \dim_{\text{H}} \mathcal{C} + \epsilon$ .*

*Proof.* Let  $E_n$  be as in Theorem 2.1 with  $n$  large enough so that  $\text{cdim}_{\text{H}} E_n < 1 - \dim_{\text{H}} \mathcal{C} + \epsilon$ , and let  $r \in [0, 2]$ . Then  $r' = 2 - r \in [0, 2]$ , and there are points  $x \in \mathcal{C}$  and  $y \in E_n$  such that  $x + y = 2 - r$ . Thus  $x + r \in 2 - E_n$ ; hence

$$\text{cdim}_{\text{H}}\{x + r\} \leq \text{cdim}_{\text{H}}(-E_n + 2) = \text{cdim}_{\text{H}} E_n < 1 - \dim_{\text{H}} \mathcal{C} + \epsilon$$

as desired.  $\square$

As we shall see in Section 3, we can even achieve a closed set  $E$  of effective Hausdorff dimension  $1 - \dim_{\text{H}} \mathcal{C}$  satisfying  $\frac{1}{2}\mathcal{C} + \frac{1}{2}E = [0, 1]$ .

### 3. LOWER BOUND

In Section 2 we demonstrated the existence of points in the Cantor set which cancel randomness; we now show there are many such points. Instead of searching for individual points with small dimension, we now characterize the Hausdorff dimension (and effective Hausdorff dimension) of all such points. We use Lorentz's Lemma again to upgrade Theorem 2.1 and Corollary 2.2. Our upgrade proceeds in two phases. The second phase occurs later in Section 5 as it relies on the upper bound results from Section 4. Our procedure is the same as that used in [4].

**Definition 3.1.** The *density* of a set  $A = \{k_1 < k_2 < k_3 < \dots\} \subseteq \mathbf{N}$  is defined to be

$$\text{density}(A) = \lim_{n \rightarrow \infty} \frac{\text{card}(A \cap \{1, 2, \dots, n\})}{n},$$

provided this limit exists.

We note that  $\text{density}(A) = \lim_{n \rightarrow \infty} \frac{n}{k_n}$ . Below  $A[n]$  will denote the length  $n$  prefix of  $A$ 's characteristic function and  $\lfloor x \rfloor$  is the integer part of  $x$ .

**Theorem 3.2.** *Let  $1 - \dim_{\text{H}} \mathcal{C} \leq \alpha \leq 1$  and let  $r \in [0, 1]$ . Then*

$$\dim_{\text{H}}[(\mathcal{C} + r) \cap \mathcal{E}_{\leq \alpha}] \geq \alpha - 1 + \dim_{\text{H}} \mathcal{C}.$$

*Proof.* For  $\alpha = 1$ , the theorem clearly holds. Thus assume

$$D = \frac{1 - \alpha}{\dim_{\text{H}} \mathcal{C}} > 0,$$

and define

$$(3.1) \quad A = \{\lfloor y/D \rfloor : y \in \mathbf{N}\}$$

so that  $D = \text{density}(A)$ . Let  $\mathcal{C}_A$  ( $\mathcal{C}_{\bar{A}}$ ) be the set of  $x \in [0, 1]$  having a ternary expansion whose digits are all 0 or 2, where the 2's only occur at positions in  $A$  (positions not in  $A$ ).

Now

$$\overline{\dim}_{\text{B}} \mathcal{C}_A \leq \limsup_{n \rightarrow \infty} \frac{\log 2^n}{\log 3^{k_n - 1}} = \dim_{\text{H}} \mathcal{C} \cdot \limsup_{n \rightarrow \infty} \frac{n}{k_n - 1} = 1 - \alpha.$$

Since upper box counting dimension dominates Hausdorff dimension [6, p. 43], we also have  $\dim_{\text{H}} \mathcal{C}_A \leq \overline{\dim}_{\text{B}} \mathcal{C}_A \leq 1 - \alpha$ . As in (2.4), the Lipschitz map  $(x, y) \mapsto x + y$  does not increase dimension [5, Lemma 1.8], so  $\dim_{\text{H}}(\mathcal{C}_A + \mathcal{C}_{\bar{A}}) \leq \dim_{\text{H}}(\mathcal{C}_A \times \mathcal{C}_{\bar{A}})$ . Since  $\mathcal{C} = \mathcal{C}_A + \mathcal{C}_{\bar{A}}$ , it follows from [6, Product formula 7.3] that

$$(3.2) \quad \dim_{\text{H}} \mathcal{C}_{\bar{A}} \geq \dim_{\text{H}} \mathcal{C} - \overline{\dim}_{\text{B}} \mathcal{C}_A \geq \dim_{\text{H}} \mathcal{C} + \alpha - 1.$$

We pause from the main argument to prove the following two lemmas. First we exploit the special form of the set  $A$ .

**Lemma 3.3.**  $K(A[n]) \leq 4 \log_3 n + O(1)$ .

*Proof of Lemma 3.3.* Let  $\frac{r}{s}$  be the largest fraction with  $s \leq n$  such that  $\frac{r}{s} \leq \frac{1}{D}$ . Notice if we know  $r$ ,  $s$ , and  $n$ , we can compute  $A[n]$  because

$$\left\lfloor \frac{y}{D} \right\rfloor = \left\lfloor \frac{ry}{s} \right\rfloor$$

for  $1 \leq y \leq n$ . (To see this, notice that if  $x = \lfloor \frac{y}{D} \rfloor > \frac{ry}{s}$ , then  $\frac{ry}{s} < x \leq \frac{y}{D}$ . This would give us  $\frac{r}{s} < \frac{x}{y} \leq \frac{1}{D}$ , contradicting maximality of  $\frac{r}{s}$ .) Specifying  $r$ ,  $s$ , and  $n$  requires a ternary string of length at most

$$(3.3) \quad \log_3 \left( \frac{n}{D} \right) + \log_3 n + \log_3 n + (2 \log_3 \log_3 n + 1) + O(1),$$

where the “ $2 \log_3 \log_3 n + 1$ ” bits are used to mark the ends of the “ $\log_3 n$ ” bit strings, and  $O(1)$  tells the universal machine how to process the input. The lemma now follows by noting that  $4 \log_3 n + O(1)$  is an upper bound for (3.3).  $\square$

**Lemma 3.4.** *There exists a closed set  $E$  such that  $\text{cdim}_{\mathbb{H}} E \leq \alpha$  and  $\mathcal{C}_A + E = [0, 2]$ .*

*Proof of Lemma 3.4.* We follow the outline of our prior argument from Section 2. The idea is to take  $E$  to be a set generated by concatenating elements from the blocks  $B_1, B_2, B_3, \dots$  as in Theorem 2.1.

For each  $k > 0$ , let  $m_k = k^2$ , let  $n_k$  denote the difference  $m_k - m_{k-1}$ , and let

$$I_k = \{i : 0 \leq i < 3^{n_k}\}.$$

Let  $A$  be the set from (3.1), and define

$$C_k = \{i \in I_k : i \text{ has only 0's and 1's in its ternary expansion} \\ \text{and the 1's only occur at positions in } A - m_{k-1}\}.$$

By Lorentz's Lemma, there exists a set  $B_k \subseteq (-3^{n_k}, 3^{n_k})$  with

$$(3.4) \quad I_k \subseteq C_k + B_k$$

satisfying, for all  $\epsilon > 0$  and all sufficiently large  $k$ ,

$$\text{card } B_k \leq c' \cdot 3^{n_k} \cdot \frac{\log [2^{n_k(D+\epsilon)}]}{2^{n_k[D-\epsilon]}} = c \cdot 3^{n_k} \cdot \frac{n_k(D+\epsilon)}{2^{n_k(D-\epsilon)}}$$

where  $c'$  is the constant obtained from Lorentz's Lemma,  $c = c' \log 2$ , and again  $D = \text{density}(A)$ . Let

$$\frac{1}{2}\mathcal{C}_A = \left\{ \sum_{k=1}^{\infty} \frac{a_k}{3^{m_k}} : a_k \in C_k \right\} \quad \text{and} \quad \frac{1}{2}E = \left\{ \sum_{k=1}^{\infty} \frac{b_k}{3^{m_k}} : b_k \in B_k \right\}.$$

The set  $E$  is closed since it is the countable intersection of closed sets. Let  $x \in [0, 1]$  with ternary expansion  $0.x_1x_2x_3\dots$ . By (3.4), there exist sequences  $\{a_k\}$  with members in  $C_k$  and  $\{b_k\}$  from  $B_k$  such that for all  $k$ ,  $\sum_{j=0}^{n_k-1} x_{m_k-j}3^j = a_k + b_k$ , and therefore

$$x = \sum_{k=1}^{\infty} \sum_{j=m_{k-1}+1}^{m_k} \frac{x_j}{3^j} = \sum_{k=1}^{\infty} \frac{1}{3^{m_k}} \sum_{j=0}^{n_k-1} x_{m_k-j}3^j = \sum_{k=1}^{\infty} \frac{a_k}{3^{m_k}} + \sum_{k=1}^{\infty} \frac{b_k}{3^{m_k}}.$$

is a member of  $\frac{1}{2}\mathcal{C}_A + \frac{1}{2}E$ . This proves  $\mathcal{C}_A + E = [0, 2]$ .



It remains to verify that  $\text{cdim}_{\mathbb{H}} E \leq \alpha$ . Let  $\epsilon > 0$ , and let  $x \in E$ . We want to compute an upper bound on  $K(x \upharpoonright m_k)$ . To specify  $x \upharpoonright m_k$ , we can first specify the sets  $B_j$ , for  $j \leq k$  and then specify which element of  $B_1 \times \cdots \times B_k$  gives the blocks of  $x \upharpoonright m_k$ .

If we know  $A[m_k]$ , we can determine the sequence of sets  $B_j$ , for  $j \leq k$  (just use a brute force search to find the first  $B_j$  as in the conclusion of Lorentz's Lemma); by Lemma 3.3 this requires a ternary string of length at most  $4 \log m_k + O(1)$  (plus an additional  $o(\log_3 m_k)$  for starting and ending delimiters if desired). An element of the known set  $B_1 \times \cdots \times B_k$  can be specified by a ternary string of length at most

$$\begin{aligned} \log_3 \prod_{j=1}^k \text{card } B_j &\leq \log_3 \prod_{j=1}^k c \cdot 3^{n_j} \cdot \frac{n_j(D+\epsilon)}{2^{n_j(D-\epsilon)}} \\ &= k \log_3 c + m_k + \log_3 \prod_{j=1}^k \frac{n_j(D+\epsilon)}{2^{n_j(D-\epsilon)}} \leq k \log_3 c + m_k + \log_3 \frac{(D+\epsilon)^k \cdot (m_k)^k}{2^{m_k \cdot (D-\epsilon)}} \\ &\leq k \log_3 c + k^2 + k \log_3(D+\epsilon) + 2k \log_3 k - k^2(D-\epsilon) \dim_{\mathbb{H}} \mathcal{C}. \end{aligned}$$

(and again we can add  $O(\log_3 m_k)$  for delimiters). Therefore,

$$\begin{aligned} K(x \upharpoonright m_k) &\leq k^2[1 - (D-\epsilon) \dim_{\mathbb{H}} \mathcal{C} + o(1)] + 4 \log_3 m_k + O(\log_3 m_k) \\ &= k^2[\alpha + \epsilon \cdot \dim_{\mathbb{H}} \mathcal{C} + o(1)], \end{aligned}$$

and appealing to the Kolmogorov complexity definition for constructive dimension (1.5), we find

$$\text{cdim}\{x\} \leq \liminf_{k \rightarrow \infty} \frac{K(x \upharpoonright m_k)}{m_k} \leq \alpha + \epsilon \cdot \dim_{\mathbb{H}} \mathcal{C}.$$

It follows from (1.4) that  $\text{cdim}_{\mathbb{H}} E \leq \alpha + \epsilon$  for every  $\epsilon > 0$ .  $\square$

Take  $E$  as in Lemma 3.4 and let  $F = 2 - E$ . Then  $F \subseteq \mathcal{E}_{\leq \alpha}$  and  $F - \mathcal{C}_A = 2 - (E + \mathcal{C}_A) = [0, 2]$ . Fix  $r \in [0, 1]$  and let  $S = \mathcal{C} \cap (F - r)$ ; it will suffice to show that  $\dim_{\mathbb{H}} S \geq \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C}$ .

Now for each  $z \in \mathcal{C}$  there exist unique points  $v \in \mathcal{C}_A$  and  $w \in \mathcal{C}_{\bar{A}}$  such that  $v + w = z$ ; let  $p$  be the projection map which takes  $z \in \mathcal{C}$  to its unique counterpart  $w \in \mathcal{C}_{\bar{A}}$ . For each  $y \in \mathcal{C}_{\bar{A}}$  we have  $r + y \in [0, 2] \subseteq F - \mathcal{C}_A$ , so there exists  $x \in \mathcal{C}_A$  such that  $r + y \in F - x$ , which gives  $x + y \in S$  since  $\mathcal{C}_A + \mathcal{C}_{\bar{A}} = \mathcal{C}$ . Thus  $p$  maps  $S$  onto  $\mathcal{C}_{\bar{A}}$ . Since  $p$  is Lipschitz we have, using (3.2),

$$(3.5) \quad \dim_{\mathbb{H}} S \geq \dim_{\mathbb{H}} \mathcal{C}_{\bar{A}} \geq \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C}$$

because Lipschitz maps do not increase dimension [5, Lemma 1.8]. Theorem 3.2 follows.  $\square$

*Remark.* The set  $E$  constructed in Lemma 3.4 has both Hausdorff dimension and effective Hausdorff dimension  $\alpha$ . Following the method of (3.2), we can establish the following lower bound:

$$\dim_{\mathbb{H}} E \geq \dim_{\mathbb{H}}[0, 2] - \overline{\dim}_{\mathbb{B}} \mathcal{C}_A \geq 1 + \alpha - 1 = \alpha.$$

## 4. UPPER BOUND

In this section we prove the following upper bound which matches the lower bound of Theorem 3.2 and Theorem 5.1.

**Theorem 4.1.** *Let  $1 - \dim_{\mathbb{H}} \mathcal{C} \leq \alpha \leq 1$ . For every Martin-Löf random real  $r$ ,*

$$(4.1) \quad \dim_{\mathbb{H}} [(\mathcal{C} + r) \cap \mathcal{E}_{\leq \alpha}] \leq \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C}.$$

*Proof.* The case  $\alpha = 1$  is trivial, so assume  $\alpha < 1$ . Fix a computable  $\gamma > \alpha + \dim_{\mathbb{H}} \mathcal{C} \geq 1$ , and let  $t = \gamma - 1$ . Let  $f$  be defined as in (1.6) and, as before, let  $B = f^{-1}(\mathcal{C} \times \mathcal{E}_{\leq \alpha})$ , so that the vertical fiber of  $B$  at  $x$  is  $B_x = (\mathcal{C} + x) \cap \mathcal{E}_{\leq \alpha}$ . To prove Theorem 4.1 it suffices to prove the following lemma.

**Lemma 4.2.** *For every Martin-Löf random real  $r$ ,  $\mathcal{H}^t(B_r) = 0$ .*

Let  $\mathcal{M}^t$  be the  $t$ -dimensional net measure in the plane induced by the net of standard dyadic squares, and for each  $\delta > 0$ , let  $\mathcal{M}_{\delta}^t$  be the  $\delta$ -approximate net measure [5].  $\mathcal{M}^t$  and  $\mathcal{M}_{\delta}^t$  are defined in the same way as  $\mathcal{H}^t$  and  $\mathcal{H}_{\delta}^t$  except that the covers  $\mathcal{G}$  from the definition in (1.3) consist exclusively of square sets of the form

$$\left[ \frac{m}{2^k}, \frac{m+1}{2^k} \right) \times \left[ \frac{n}{2^k}, \frac{n+1}{2^k} \right)$$

for integers  $k$ ,  $m$ , and  $n$ . Hence for any set  $E$ ,  $\mathcal{M}_{\delta}^t(E) \geq \mathcal{H}_{\delta}^t(E)$ . Let  $\mathcal{L}^1 = \mathcal{H}^1$  be Lebesgue measure on the real line.

To prove Lemma 4.2, it suffices to show the following which is a “computable” version of Marstrand’s method below. Indeed any  $x$  which belongs to the left-hand side of (4.2) for every  $m$  fails a Martin-Löf test and therefore cannot be Martin-Löf random.

**Lemma 4.3.** *Fix  $c > 0$ . There is a uniformly computable sequence of open sets  $U_m$  with  $\mathcal{L}^1(U_m) < 2^{-m}$  such that for each  $m$ ,*

$$(4.2) \quad \{x : \mathcal{H}^t(B_x) > c\} \subseteq U_m.$$

*Proof of Lemma 4.3.* We may assume that  $c$  is computable. Fix a uniformly computable sequence of collections  $\mathcal{S}_k = \{S_{k,i}\}$  of dyadic squares which for each  $k$ , forms a  $2^{1/2} \cdot \frac{1}{k}$ -mesh cover of  $B$  with

$$\sum_{S_{k,i} \in \mathcal{S}_k} |S_{k,i}|^{\gamma} < \frac{c \cdot 2^{1/2}}{2^{k+1}}.$$

We show the existence of such a sequence in Lemma 4.5. For each  $k$ , let

$$A_k = \left\{ x : \sum_{i=0}^{\infty} |(S_{k,i})_x|^t > c \right\}$$

where  $(S_{k,i})_x$  denotes the vertical fiber of  $S_{k,i}$  at  $x$ . The sets  $A_k$  are unions of left-closed right-open dyadic intervals in a uniformly computable way. Since

$$\mathcal{H}^t(B_x) > c \implies x \in \bigcap_{m=1}^{\infty} \bigcup_{k=m}^{\infty} A_k,$$

we shall see that it suffices to show  $\mathcal{L}^1(A_k) < 2^{-k-1}$ .

Let  $a_{k,i} = |S_{k,i}|^t$  and let

$$I_{k,i} = \{x : (x, y) \in S_{k,i} \text{ for some } y\}.$$

Each  $I_{k,i}$  is a dyadic interval and  $|I_{k,i}| = 2^{-1/2}|S_{k,i}|$ . Also, if  $x \in A_k$ , then  $\sum_{\{i:x \in I_{k,i}\}} a_{k,i} > c$ . We now appeal to [5, Lemma 5.7]:

**Marstrand's Lemma** ([11]). *Let  $A \subseteq \mathbf{R}$ , let  $\{I_n\}$  be a  $\delta$ -mesh cover of  $A$  by dyadic intervals, and let  $a_n > 0$  for all  $n$ . Suppose that for all  $x \in A$*

$$\sum_{\{n:x \in I_n\}} a_n > c$$

for some constant  $c$ . Then for all  $s$ ,

$$\sum_n a_n |I_n|^s \geq c \cdot \mathcal{M}_\delta^s(A).$$

Applying Marstrand's Lemma with  $s = 1$ , we have

$$\frac{c}{2^{k+1}} > 2^{-1/2} \sum_{S_{k,i} \in \mathcal{S}_k} |S_{k,i}|^\gamma = \sum_{\{i:x \in I_{k,i}\}} a_{k,i} |I_{k,i}| \geq c \cdot \mathcal{M}_{1/k}^1(A_k).$$

It follows that  $\mathcal{L}^1(A_k) < 2^{-k-1}$ . Thus, for each  $m$ ,  $\mathcal{L}^1(\cup_{k=m}^\infty A_k) < 2^{-m}$ . Now we may find uniformly computable open sets  $U_m$  with

$$\{x : \mathcal{H}^t(B_x) > c\} \subseteq \bigcup_{k=m}^\infty A_k \subseteq U_m$$

and  $\mathcal{L}^1(U_m) < 2^{-m}$ . □

For  $s \in [0, 1]$ , a *weak  $s$ -randomness test* [17] is a sequence of uniformly c.e. sets of open dyadic intervals  $U_0, U_1, U_2, \dots$  such that  $\sum_{\sigma \in U_n} 2^{-s|\sigma|} \leq 2^{-n}$  for all  $n$ . We will call a set  $E \subseteq \mathbf{R}$  *weakly  $s$ -random* if  $E \not\subseteq \bigcap_n U_n$  for every weak  $s$ -randomness test  $U_0, U_1, U_2, \dots$ . We will need the following technical result in order to ensure that the cover  $\mathcal{S}_k$  in Lemma 4.5 is sufficiently uniform:

**Lemma 4.4.** *For every  $d > \alpha$  there is a computable function  $\langle j, k, l \rangle \mapsto Q_{\langle k, l \rangle, j}$  such that for all  $k$  and  $l$ ,*

- (I)  $\{Q_{\langle k, l \rangle, j}\}_j$  is a  $2^{-k}$ -mesh cover of  $\mathcal{E}_{\leq \alpha}$  by dyadic intervals, and
- (II)  $\sum_{j=0}^\infty |Q_{\langle k, l \rangle, j}|^d < 2^{-l}$ .

*Proof of Lemma 4.4.* Without loss of generality, we can assume that  $l \geq kd$ ; proving the result for a larger  $l$  only makes the second part of the lemma more true. An inspection of [3, Proposition 13.5.3] (and the proof of the theorem immediately preceding it) reveals that for any  $s \in [0, 1]$  and any set of reals  $E$ ,

$$\text{cdim } E \geq \sup\{s : E \text{ is weakly } s\text{-random}\}.$$

Since  $\text{cdim}_H(\mathcal{E}_{\leq \alpha}) = \alpha < d$  [10, Theorem 4.7], we have that  $\mathcal{E}_{\leq \alpha}$  is not weakly  $d$ -random. This means that there exists a uniformly c.e. collection of dyadic intervals  $Q_{k,l} = \{Q_{\langle k, l \rangle, j} : j \geq 0\}$  such that

$$(4.3) \quad \sum_{j=0}^\infty |Q_{\langle k, l \rangle, j}|^d \leq 2^{-l}$$

and each  $\mathcal{Q}_{k,l}$  covers  $\mathcal{E}_{\leq\alpha}$  which proves (II). It follows from (4.3) that for every  $j_0$ ,  $|Q_{\langle k,l \rangle, j_0}|^d \leq 2^{-l}$ , and so

$$|Q_{\langle k,l \rangle, j_0}| \leq 2^{-l/d} \leq 2^{-k}$$

as needed for (I).  $\square$

**Lemma 4.5.** *There exists a uniformly computable sequence of collections of dyadic squares  $\mathcal{S}_k$  which, for each  $k$ , form a  $2^{1/2} \cdot \frac{1}{k}$ -mesh cover of  $B$  with*

$$\sum_{S \in \mathcal{S}_k} |S|^\gamma < \frac{c \cdot 2^{1/2}}{2^{k+1}}.$$

*Proof of Lemma 4.5.* Write  $\gamma = s + d$ , where  $s > \dim_{\mathbb{H}} \mathcal{C}$  and  $d > \alpha$ . Let  $\mathcal{G}_k$  be a uniformly computable mesh cover of  $\mathcal{C}$  by dyadic intervals of length  $2^{-k}$  such that for each  $k$ ,  $\text{card}(\mathcal{G}_k) \leq 2^{sk}$  and, let  $\mathcal{Q}_{k,l} = \{Q_{\langle k,l \rangle, j} : j \geq 0\}$  as in Lemma 4.4. Form the uniformly computable sequence of square covers:

$$\Gamma_{k,l} = \{G \times Q : Q \in \mathcal{Q}_{k,l} \text{ and } G \in \mathcal{G}_{-\log |Q|}\}.$$

Then, using  $\text{card}[\mathcal{G}_{-\log |Q|}] \leq |Q|^{-s}$  for all  $Q \in \mathcal{Q}_{k,l}$ ,

$$\begin{aligned} \sum_{X \in \Gamma_{k,l}} |X|^\gamma &= \sum_{\substack{Q \in \mathcal{Q}_{k,l} \\ G \in \mathcal{G}_{-\log |Q|}}} |G \times Q|^\gamma = 2^{\gamma/2} \sum_{Q \in \mathcal{Q}_{k,l}} |Q|^{s+d} \cdot \text{card}[\mathcal{G}_{-\log |Q|}] \\ &\leq 2^{\gamma/2} \sum_{Q \in \mathcal{Q}_{k,l}} |Q|^d < 2^{-l}. \end{aligned}$$

Let  $f$  be the Lipschitz mapping (1.6) whose inverse map does not increase diameter by more than a factor of  $\sqrt{2}$  and maps  $\Gamma_{k,l}$  onto  $B$  for all  $k$  and  $l$ . For every  $k$ , let  $m(k)$  be sufficiently large so that  $2^{-m(k)}$  is less than  $c/2^{k+1}$ . Now form the collection  $\mathcal{S}_k$  by taking, for each  $X \in \Gamma_{k,m(k)}$ , the two dyadic squares which together cover the sheared dyadic square  $f^{-1}(X)$ . Then the  $\mathcal{S}_k$ 's form a uniformly computable sequence of square covers which achieves the desired bounds.  $\square$

This concludes the proof of Theorem 4.1.  $\square$

*Remark.* In contrast to the lower bound in Theorem 3.2 which holds for *all* reals in  $[0, 1]$ , the upper bound in Theorem 4.1 indeed requires some hypothesis on  $r$ . Indeed if  $r = 0$  and  $\dim_{\mathbb{H}} \mathcal{C} < \alpha < 1$  satisfied (4.1), we would have

$$\dim_{\mathbb{H}} \mathcal{C} = \dim_{\mathbb{H}} [(\mathcal{C} + 0) \cap \mathcal{E}_{\leq\alpha}] \leq \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C} < \dim_{\mathbb{H}} \mathcal{C},$$

a contradiction.

## 5. LOWER BOUND II

We modify the proof of Theorem 3.2 to obtain a stronger result for the case of Martin-Löf randoms:

**Theorem 5.1.** *Let  $1 - \dim_{\mathbb{H}} \mathcal{C} \leq \alpha \leq 1$  and let  $r \in [0, 1]$  be Martin-Löf random. Then*

$$\dim_{\mathbb{H}} [(\mathcal{C} + r) \cap \mathcal{E}_{=\alpha}] = \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C}.$$

*Moreover,*

$$\mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}} [(\mathcal{C} + r) \cap \mathcal{E}_{=\alpha}] > 0.$$

*Proof.* Fix an  $\alpha$  satisfying  $1 - \dim_{\mathbb{H}} \mathcal{C} \leq \alpha \leq 1$ , and let  $A, \mathcal{C}_A, \mathcal{C}_{\bar{A}}, E, F$ , and  $S$  be as in the proofs of Theorem 3.2 and Lemma 3.4. For  $x \in \mathbf{R}$ , let

$$N_{\delta}(x) = \{y \in \mathbf{R} : |x - y| \leq \delta\}.$$

We shall make use of the following result from Mattila's book:

**Theorem 5.2** ([13], Theorem 6.9). *Let  $\mu$  be a Radon measure on  $\mathbf{R}^n$ ,  $E \subseteq \mathbf{R}^n$ ,  $0 < \lambda < \infty$ , and  $\alpha > 0$ . If*

$$\limsup_{\delta \rightarrow 0} \frac{\mu[N_{\delta}(x)]}{(2\delta)^{\alpha}} \leq \lambda$$

for all  $x \in E$ , then  $\mathcal{H}^{\alpha}(E) \geq \frac{\mu(E)}{2^{\alpha}\lambda}$ .

**Lemma 5.3.**  $\mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}(\mathcal{C}_{\bar{A}}) > 0$ .

*Proof of Lemma 5.3.* Since the case  $\alpha = 1 - \dim_{\mathbb{H}} \mathcal{C}$  is trivial, we assume  $\alpha > 1 - \dim_{\mathbb{H}} \mathcal{C}$ . Let  $\beta = \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C}$ . Since  $A = \{\lfloor y/D \rfloor : y \in \mathbf{N}\}$  where  $D = \frac{1-\alpha}{\dim_{\mathbb{H}} \mathcal{C}}$ , we have  $\bar{A} = \{u_1 < u_2 < u_3 < \dots\}$ , where  $\lim_{n \rightarrow \infty} \frac{u_n}{n} = 1 - D$ . In fact, the careful choice of the set  $A$  lets us make a stronger statement about the numbers  $u_n$ : there is a fixed number  $t$  such that  $u_n \leq (n+t)/(1-D)$  for all  $n$ . For each finite binary string  $\sigma = \sigma_1\sigma_2 \dots \sigma_n$ , let  $I(\sigma)$  be the closed interval of ternary expansions  $x = 0.x_1x_2 \dots$  satisfying

$$x_p = \begin{cases} \sigma_k & \text{if } p = u_k \text{ for some } k \leq n, \text{ and} \\ 0 & \text{if } p \leq u_n \text{ and } p \notin \{u_1, \dots, u_n\}. \end{cases}$$

Define a probability measure  $\mu$  on  $[0, 1]$  by requiring

$$\mu[I(\sigma) \cap \mathcal{C}_{\bar{A}}] = \frac{1}{2^{|\text{length of } \sigma|}}.$$

Since  $\mu$  is a bounded Borel measure supported on the compact set  $\mathcal{C}_{\bar{A}}$ ,  $\mu$  is a Radon measure; hence Theorem 5.2 applies. For  $\delta > 0$ , let  $f(\delta)$  be the least index such that  $\delta > 3^{-u_{f(\delta)}}$ . Let

$$x = .000 \dots 00x_{u_1}000 \dots 00x_{u_2}000 \dots 00x_{u_{f(\delta)}}000 \dots \in \frac{1}{2}\mathcal{C}_{\bar{A}},$$

and let  $J = I[x_{u_1}x_{u_2} \dots x_{u_{f(\delta)}}]$ . Then  $x \in J$  and the length of the closed interval  $J$  is  $3^{-u_{f(\delta)}} < \delta$ . So  $J \subseteq N_{\delta}(x)$ . Now the length of each interval  $I[\sigma_1 \dots \sigma_{f(\delta)-1}]$  is  $3^{-u_{f(\delta)-1}} \geq \delta$ , so  $N_{\delta}(x)$  can intersect no more than 4 of these non-overlapping intervals. Therefore  $\mu[N_{\delta}(x)] \leq 4 \cdot 2^{-(f(\delta)-1)}$ . Let  $c = 1 - D$ . Then

$$\frac{\mu[N_{\delta}(x)]}{(2\delta)^{\beta}} \leq \frac{4 \cdot 2^{-(f(\delta)-1)}}{(2 \cdot 3^{-u_{f(\delta)}})^{\beta}} \leq \frac{8}{2^{\beta}} \cdot \left( \frac{3^{\beta u_{f(\delta)}}}{2^{f(\delta)}} \right) \leq \frac{8}{2^{\beta}} \cdot \left( \frac{3^{\beta \cdot \frac{f(\delta)+t}{c}}}{2^{f(\delta)}} \right).$$

Thus

$$\limsup_{\delta \rightarrow 0} \frac{\mu[N_{\delta}(x)]}{(2\delta)^{\beta}} \leq \limsup_{\delta \rightarrow 0} 8 \cdot \frac{3^{\beta t/c}}{2^{\beta}} \cdot \left( \frac{3^{\beta/c}}{2} \right)^{f(\delta)} = 8 \cdot \frac{3^{\beta t/c}}{2^{\beta}},$$

and hence by Theorem 5.2, we have  $\mathcal{H}^{\beta}(\frac{1}{2}\mathcal{C}_{\bar{A}}) \geq \frac{3^{-\beta t/c}}{8}$ . It follows that  $\mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}(\mathcal{C}_{\bar{A}}) = \mathcal{H}^{\beta}(\mathcal{C}_{\bar{A}}) > 0$ .  $\square$

**Lemma 5.4.**  $\mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}[(\mathcal{C} + r) \cap \mathcal{E}_{\leq \alpha}] > 0$ .

*Proof of Lemma 5.4.* Let  $S = \mathcal{C} \cap (E - r)$ . By Lemma 3.4,  $E \subseteq \mathcal{E}_{<\alpha}$ , and so it suffices to show that  $\mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}(S) > 0$ . Retracing the argument of Theorem 3.2 down to (3.5), we get  $\dim_{\mathbb{H}} S \geq \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C}$ . Furthermore, as we now argue,

$$(5.1) \quad \mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}(S) \geq q_{\alpha} \cdot \mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}(\mathcal{C}_A) > 0$$

for some constant  $q_{\alpha} > 0$ . The strict inequality in (5.1) follows from Lemma 5.3. For the nonstrict inequality, we again appeal to [5, Lemma 1.8] and the fact that the projection map from  $\mathcal{C}$  to  $\mathcal{C}_A$  is Lipschitz. [5, Lemma 1.8] states that, up to some constant factor, a Lipschitz map does not decrease  $\mathcal{H}^{\alpha}$  measure. This is slightly stronger than what we used before in Theorem 3.2, namely that a Lipschitz map cannot increase dimension.  $\square$

Using the assumption that  $r$  is Martin-Löf random, we next obtain the following:

**Lemma 5.5.**  $\mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}[(\mathcal{C} + r) \cap \mathcal{E}_{<\alpha}] = 0$ .

*Proof of Lemma 5.5.* The case  $\alpha = 1 - \dim_{\mathbb{H}} \mathcal{C}$  follows from Theorem 1.1, so assume  $\alpha > 1 - \dim_{\mathbb{H}} \mathcal{C}$ . By Theorem 4.1, for any  $\gamma$  with  $1 - \dim_{\mathbb{H}} \mathcal{C} \leq \gamma < \alpha$ , we have

$$\dim_{\mathbb{H}}[(\mathcal{C} + r) \cap \mathcal{E}_{\leq \gamma}] \leq \gamma - 1 + \dim_{\mathbb{H}} \mathcal{C} < \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C}.$$

Since  $\mathcal{E}_{<\alpha}$  is the countable union of sets  $E_{\leq \gamma}$  for a sequence of  $\gamma$ 's approaching  $\alpha$  from below, the theorem follows.  $\square$

Combining Lemma 5.4 with Lemma 5.5, we find that

$$\mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}[(\mathcal{C} + r) \cap \mathcal{E}_{=\alpha}] = \mathcal{H}^{\alpha-1+\dim_{\mathbb{H}} \mathcal{C}}[(\mathcal{C} + r) \cap (\mathcal{E}_{\leq \alpha} - \mathcal{E}_{<\alpha})] > 0,$$

whence we conclude the desired theorem.  $\square$

Combining Theorem 5.1 with Theorem 3.2 and Theorem 4.1 yields our final result:

**Corollary 5.6.** *For any  $\alpha$  satisfying  $1 - \dim_{\mathbb{H}} \mathcal{C} \leq \alpha \leq 1$ , and for any Martin-Löf random  $r \in [0, 1]$ , we have*

$$\dim_{\mathbb{H}}[(\mathcal{C} + r) \cap \mathcal{E}_{=\alpha}] = \dim_{\mathbb{H}}[(\mathcal{C} + r) \cap \mathcal{E}_{\leq \alpha}] = \alpha - 1 + \dim_{\mathbb{H}} \mathcal{C}.$$

#### REFERENCES

- [1] Krishna B. Athreya, John M. Hitchcock, Jack H. Lutz, and Elvira Mayordomo. Effective strong dimension in algorithmic information and computational complexity. *SIAM J. Comput.*, 37(3):671–705 (electronic), 2007. ISSN 0097-5397. doi: 10.1137/S0097539703446912. URL <http://dx.doi.org/10.1137/S0097539703446912>.
- [2] Jin-yi Cai and Juris Hartmanis. On Hausdorff and topological dimensions of the Kolmogorov complexity of the real line. *Journal of Computer and System Sciences*, 49(3):605–619, 1994. ISSN 0022-0000. doi: DOI:10.1016/S0022-0000(05)80073-X. 30th IEEE Conference on Foundations of Computer Science.
- [3] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic randomness and complexity*. Theory and Applications of Computability. Springer, New York, 2010. ISBN 978-0-387-95567-4.
- [4] P. Erdős, K. Kunen, and R. Daniel Mauldin. Some additive properties of sets of real numbers. *Fund. Math.*, 113(3):187–199, 1981. ISSN 0016-2736.

- [5] K. J. Falconer. *The geometry of fractal sets*, volume 85 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1986. ISBN 0-521-25694-1; 0-521-33705-4.
- [6] Kenneth Falconer. *Fractal geometry*. John Wiley & Sons Inc., Hoboken, NJ, second edition, 2003. ISBN 0-470-84861-8. doi: 10.1002/0470013850. URL <http://dx.doi.org/10.1002/0470013850>. Mathematical foundations and applications.
- [7] L. A. Levin. The concept of a random sequence. *Dokl. Akad. Nauk SSSR*, 212: 548–550, 1973. ISSN 0002-3264.
- [8] Ming Li and Paul Vitányi. *An introduction to Kolmogorov complexity and its applications*. Texts in Computer Science. Springer, New York, third edition, 2008. ISBN 978-0-387-33998-6. doi: 10.1007/978-0-387-49820-1. URL <http://dx.doi.org/10.1007/978-0-387-49820-1>.
- [9] G. G. Lorentz. On a problem of additive number theory. *Proc. Amer. Math. Soc.*, 5:838–841, 1954. ISSN 0002-9939.
- [10] Jack H. Lutz. The dimensions of individual strings and sequences. *Inform. and Comput.*, 187(1):49–79, 2003. ISSN 0890-5401. doi: 10.1016/S0890-5401(03)00187-1. URL [http://dx.doi.org/10.1016/S0890-5401\(03\)00187-1](http://dx.doi.org/10.1016/S0890-5401(03)00187-1).
- [11] J. M. Marstrand. The dimension of Cartesian product sets. *Proc. Cambridge Philos. Soc.*, 50:198–202, 1954.
- [12] Per Martin-Löf. The definition of random sequences. *Information and Control*, 9:602–619, 1966. ISSN 0890-5401.
- [13] Pertti Mattila. *Geometry of sets and measures in Euclidean spaces*, volume 44 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1995. ISBN 0-521-46576-1; 0-521-65595-1. doi: 10.1017/CBO9780511623813. URL <http://dx.doi.org/10.1017/CBO9780511623813>. Fractals and rectifiability.
- [14] Elvira Mayordomo. A Kolmogorov complexity characterization of constructive Hausdorff dimension. *Inform. Process. Lett.*, 84(1):1–3, 2002. ISSN 0020-0190. doi: 10.1016/S0020-0190(02)00343-5. URL [http://dx.doi.org/10.1016/S0020-0190\(02\)00343-5](http://dx.doi.org/10.1016/S0020-0190(02)00343-5).
- [15] Jan Reimann. *Computability and Fractal Dimension*. PhD thesis, University of Heidelberg, December 2004.
- [16] Robert I. Soare. *Recursively enumerable sets and degrees*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1987. ISBN 3-540-15299-7. A study of computable functions and computably generated sets.
- [17] Kohtaro Tadaki. A generalization of Chaitin’s halting probability  $\Omega$  and halting self-similar sets. *Hokkaido Math. J.*, 31(1):219–253, 2002. ISSN 0385-4035.
- [18] William P. Ziemer. *Modern Real Analysis*. Second edition, 2004. URL <http://www.indiana.edu/~mathwz/PRbook.pdf>.

CCR-LA JOLLA, 4320 WESTERRA COURT, SAN DIEGO, CA 92121

*E-mail address:* `rdough@ccrwest.org`

IOWA STATE UNIVERSITY

*E-mail address:* `lutz@cs.iastate.edu`

UNIVERSITY OF NORTH TEXAS, DENTON, TX 76203

*E-mail address:* `mauldin@unt.edu`

RUPRECHT-KARLS-UNIVERSITÄT HEIDELBERG

*E-mail address:* `deutsch@math.uni-heidelberg.de`